

Evidence Pipeline

James Thomas



LSR Symposium





Evidence Pipeline

Finding and classifying relevant research





The new CRS-Web

CRS Web (online) | crsdemo.metaxis.com/index.php#Search

Search Simple MeSH Classifier Saved Tracking

Classifier search

Records that have been through the classifier have probabilities assigned to them to indicate how likely they are to have certain properties, like whether they are of interest to a review group, or whether they are likely to be an RCT. Choose the classifier model you are interested in, set the model parameters and click Search to find the records

RCT

Approximately 32129 records that are between 99 and 100 percent likely to be of interest

[Search](#)

You can find your records that are currently being processed by the classifier by searching for INPROCESS:CLASSIFIER

[Find those records](#)

Cochrane Register of Studies

Anna Noel-Storr [DEMENTIA] | logout

Dashboard Records Import Journals CT.GOV Reports To do

Search Layout1 Layout2 Layout3 Layout4

Search results (399 records) Page 1 of 8

#	Title	Author
1	Cognitive effects of treating obstructive sleep apnea in Alzheimer's disease: a randomized controlled study	Ancoli-Israel S // Palmer BW // Cooke
2	Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: a random...	Erkinjuntti T // Kurz A // Gauthier S //
3	Donepezil improved memory in multiple sclerosis in a randomized clinical trial	Krupp LB // Christodoulou C // Melvil
4	A randomized, 26-week, double-blind, placebo-controlled trial to evaluate the safety and efficacy of galantamine in the treatme...	Auchus A
5	A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. Donepezil Study Group	Rogers SL // Farlow MR // Doody RS /
6	A Controlled, Double-Blind, Randomized Pilot Clinical Trial of Hydroxysafflor Yellow a on Cognitive Function in Patients With Vas...	Tian J

Record

Fields Duplicates Links Reviews Classifier Files Audit CENTRAL REGISTER

The bar chart below shows the classifier scores for this record. Scores are presented in the range 0 - 100 where higher scores mean a higher likelihood that the record is of interest to the group. You can tell a group about this record if it doesn't already have it in its segment by clicking the bar for that group.

In register
 In segment
 Not in segment
 Not relevant to my group

There is a 99% likelihood that this record is an RCT [Confirm this is not an RCT] [Confirm this is an RCT]

Collection: Direct oral anticoagulants for thromboprophylaxis after orthopedic surgery

Owner: dperezrada@gmail.com

Update recommendations

Delete Collection

Related (26) Included (119) Excluded (59) Machine Recommendations (24)

Primary Study

Single intravenous administration of TB-402 for the prophylaxis of venous thromboembolism after total knee replacement: a dose-escalating, randomized, controlled trial.

AUTHORS » Verhamme P, Tangelder M, Verhaeghe R, Agenc W, Glazer S, Prins M, Jacquemin M, Büller H, TB-402 Study Group

JOURNAL » Journal of thrombosis and haemostasis : JTH

YEAR » 2011

LINKS » Pubmed, DOI

This article is included in 1 Systematic Review

Include Exclude

Systematic Review

Rivaroxaban for thromboprophylaxis after orthopaedic surgery: pooled analysis of two studies.

AUTHORS » Fisher WD, Eriksson BI, Bauer KA, Borris L, Dahl OE, Gent M, Haas S, Homering M, Huisman MV, Kakkar AK, Kälebo P, Kwong LM, Misselwitz F, Turpie AG

JOURNAL » Thrombosis and haemostasis

YEAR » 2007

LINKS » Pubmed

Without references

Include Exclude

Unclassified

Fondaparinux for prevention of venous thromboembolism in major orthopedic surgery.

AUTHORS » Tran AH, Lee G

JOURNAL » The Annals of pharmacotherapy

YEAR » 2003

LINKS » Pubmed, DOI

Include Exclude

Primary Study

Evidence of venous thromboembolism after lower limb

Abstract

About this article

Despite current guidelines recommendations about anticoagulant prophylaxis, many studies have shown an high venous thromboembolism (VTE) incidence in patients undergoing total hip and knee arthroplasty. A number of anticoagulants are currently available, but they have some limitations that affect their applicability and consequently their effectiveness. Several new oral anticoagulants (NOACs) have been developed in an attempt to overcome these limitations. Apixaban is a NOAC that selectively inhibits the coagulation factor Xa; it is approved for the prevention of VTE after total hip replacement and total knee replacement surgery. This review examines the results of main trials designed to test efficacy and safety of apixaban in major elective orthopedic surgery.



Project Transform



You can make a difference

Become a Cochrane citizen scientist. Anyone can join our collaborative volunteer effort to help categorise and summarise healthcare evidence so that we can make better healthcare decisions.

Give it a try

Cochrane Crowd
crowd.cochrane.org

Thank you

james.thomas@ucl.ac.uk

LSR Symposium



Trusted evidence.
Informed decisions.
Better health.